


Light Hadrons and New Enhancements in J/ψ Decays at BESII

Guofa XU

(for the BES Collaboration)

Institute of High Energy Physics, CAS, Beijing 100049, China



Based on 58 million J/ψ samples collected by the BESII detector at the BEPC, many mesons, baryons, and new resonances have been reported. Here, I will review some recent results of glueball candidates and new enhancement.

1 Introduction

In this paper, some recent BESII results are reported based on 58 million J/ψ events collected by the BESII detector at the BEPC. For much more detail, please see the references.

2 Scalars (0^{++})

As we know that so many scalars are listed in PDG06¹, but according to the quark model no enough room for all of these scalar particles. On the other hand, the Lattice QCD predicted that the ground state glueball is 0^{++} , and its mass is around 1.5~1.8 GeV. Theoretical physicists expect that glueballs will mix with nearby $q\bar{q}$ states of the same quantum numbers^{2,3}, it makes the situation more difficult for the glueball identification. Although the identification of a glueball is very complicated, there are several glueball candidates, such as $f_0(1500)$ and $f_0(1710)$, considering the possible mix with the ordinary $q\bar{q}$ meson, $f_0(1370)$, $f_0(1500)$, $f_0(1710)$, and $f_0(1790)$ have been analyzed for more detail by using the partial wave analyzes (PWA) method in $J/\psi \rightarrow \gamma\pi\pi$, $\gamma K\bar{K}$, $J/\psi \rightarrow \omega K\bar{K}$, and $J/\psi \rightarrow \phi\pi\pi$, $\phi K\bar{K}$ channels.

2.1 The Analysis of $J/\psi \rightarrow \gamma\pi\pi$ and $\gamma K\bar{K}$ Channels

The partial wave analyzes of $J/\psi \rightarrow \gamma\pi^+\pi^-$ and $J/\psi \rightarrow \gamma\pi^0\pi^0$ show the evidence for two 0^{++} states around the 1.45 and 1.75 GeV/ c^2 mass regions (Fig. 1, 2)⁴. The $f_0(1500)$ has

a mass of $1466 \pm 6 \pm 20 \text{ MeV}/c^2$, a width of $108_{-11}^{+14} \pm 25 \text{ MeV}/c^2$, and a branching fraction $\mathcal{B}(J/\psi \rightarrow \gamma f_0(1500) \rightarrow \gamma \pi^+ \pi^-) = (0.67 \pm 0.02 \pm 0.30) \times 10^{-4}$. The 0^{++} state in the $\sim 1.75 \text{ GeV}/c^2$ mass region has a mass of $1765_{-3}^{+4} \pm 13 \text{ MeV}/c^2$ and a width of $145 \pm 8 \pm 69 \text{ MeV}/c^2$.




Figure 1: The $\pi^+\pi^-$ invariant mass distribution from $J/\psi \rightarrow \gamma \pi^+ \pi^-$. The crosses are data, the full histogram shows the maximum likelihood fit, and the shaded histogram corresponds to the $\pi^+\pi^-\pi^0$ background.




Figure 2: The $\pi^0\pi^0$ invariant mass distribution from $J/\psi \rightarrow \gamma \pi^0 \pi^0$. The crosses are data, the full histogram shows the maximum likelihood fit, and the shaded histogram corresponds to the background.

The PWA of $J/\psi \rightarrow \gamma K^+ K^-$ and $J/\psi \rightarrow \gamma K_S^0 K_S^0$ show strong production of the $f_2'(1525)$ and the S-wave resonance $f_0(1710)$ (Fig. 3)⁵. The $f_0(1710)$ peaks at a mass of $1740 \pm 4_{-25}^{+10} \text{ MeV}$ with a width of $166_{-8}^{+5+15} \text{ MeV}$.




Figure 3: Invariant mass spectra of a) $K^+ K^-$, b) $K_S^0 K_S^0$ for $J/\psi \rightarrow \gamma K \bar{K}$ events, where the shaded histograms correspond to the estimated background contributions.

2.2 The Analysis of $J/\psi \rightarrow \omega K^+ K^-$ Channel

From Fig. 4, one can see that a dominant feature is $f_0(1710)$ ⁶. The fitted $f_0(1710)$ optimizes at $M = 1738 \pm 30 \text{ MeV}/c^2$, $\Gamma = 125 \pm 20 \text{ MeV}/c^2$.

2.3 The Analysis of $J/\psi \rightarrow \phi \pi^+ \pi^-$ and $\phi K^+ K^-$ Channels

After the partial wave analyzes for these $\phi \pi \pi$ and $\phi K K$ channels⁷, the data reported here have three important features. Firstly, the parameters of $f_0(980)$ are all well determined. Secondly,




Figure 4: (c) and (d) are projections on to K^+K^- and ωK mass. Histograms show the maximum likelihood fit; the shaded region indicates the background estimated from sidebins; the dashed curve in (d) shows the magnitude of the $K_1(1400)$ contribution and a $K\omega$ contribution at 1945 MeV/c².

there is the clearest signal to date of $f_0(1370) \rightarrow \pi^+\pi^-$; a resonant phase variation is required, from interference with $f_2(1270)$. Thirdly, there is a clear peak in $\pi\pi$ at 1775 MeV/c², consistent with $f_0(1790)$; spin 2 is less likely than spin 0.




Figure 5: The K^+K^- invariant mass distributions for (a) $J/\psi \rightarrow K^+K^-\pi^+\pi^-$, (c) $J/\psi \rightarrow K^+K^-K^+K^-$; curves show the fitted background and a Gaussian fit to the ϕ ; (b) and (d) show mass projections for events selected within ± 15 MeV/c² of the ϕ .

In summary, (1) $f_0(1370)$ has been seen in $J/\psi \rightarrow \phi\pi\pi$, but not in $J/\psi \rightarrow \omega\pi\pi$. (2) No peak of the $f_0(1500)$ directly seen in $J/\psi \rightarrow \phi KK$, ωKK , $\phi\pi\pi$, and $\omega\pi\pi$, but in proton-proton scattering is quite clear. (3) $f_0(1710)$ is observed clearly in both $J/\psi \rightarrow \phi KK$ and $J/\psi \rightarrow \omega KK$, but with $Br(J/\psi \rightarrow \omega f_0(1710) \rightarrow \omega KK)/Br(J/\psi \rightarrow \phi f_0(1710) \rightarrow \phi KK) \sim 6$, which is against a simple $s\bar{s}$ configuration for this state. (4) $f_0(1790)$ which is seen in $\pi\pi$ rather than $K\bar{K}$.

Different models have different interpretations for these experimental results. One of the interpretations is from Cheng⁸, he explained that (1) $f_0(1710)$ is composed primarily of the scalar glueball. (2) $f_0(1500)$ is close to an $SU(3)$ octet. The glueball content of $f_0(1500)$ is very tiny because an $SU(3)$ octet does not mix with the scalar glueball. (3) $f_0(1370)$ consists of an approximate $SU(3)$ singlet with some glueball component ($\sim 10\%$).

3 Pseudo-scalars (0^{-+})

The first observation of $\eta(1440)$ was made in $p\bar{p}$ annihilation at rest into $\eta(1440)\pi^+\pi^-$, $\eta(1440) \rightarrow K\bar{K}\pi$ ⁹. Nowadays, The existence of two overlapping pseudo-scalar states has been suggested to instead of the $\eta(1440)$: one around 1405 MeV/ c^2 decays mainly through $a_0(980)\pi$ (or direct $K\bar{K}\pi$), and the other around 1475 MeV/ c^2 mainly to $K^*(892)\bar{K}$ ^{1,10}. It is therefore conceivable that the higher mass state is the $s\bar{s}$ member of the 2^1S_0 nonet^{11,12}, while the lower mass state may contain a large gluonic content^{13,14}.




Figure 6: The $\gamma\rho$ invariant mass distribution. The insert shows the full mass scale where the $\eta(958)$ is clearly observed.




Figure 7: The invariant mass of $\gamma\phi$ after side-band background subtraction.

In our $J/\psi \rightarrow \gamma\gamma V$ analysis¹⁵, there is a resonance around 1424 MeV at the $J/\psi \rightarrow \gamma\gamma\rho$ channel. Comparing our result on the branching ratio $B(J/\psi \rightarrow \gamma X(1424) \rightarrow \gamma\gamma\rho) = (1.07 \pm 0.17 \pm 0.11) \times 10^{-4}$, and the upper limit of $B(J/\psi \rightarrow \gamma X(1424) \rightarrow \gamma\gamma\phi) < 0.82 \times 10^{-4}$ (95% C.L.), we cannot draw a definite conclusion on whether the $X(1424)$ is either a $q\bar{q}$ state or a glueball state.

We also analyzed the $\eta(1405)/\eta(1475)$ at $J/\psi \rightarrow \{\omega, \phi\}K\bar{K}\pi$ channels¹⁶. In the invariant mass spectra of $K_S^0 K^\pm \pi^\mp$ and $K^+ K^- \pi^0$ recoiling against the ω signal region, the resonance at 1.44 GeV/ c^2 is observed, while in the invariant mass spectra of $K_S^0 K^\pm \pi^\mp$ and $K^+ K^- \pi^0$ recoiling against the ϕ signal region, no significant structure near 1.44 GeV/ c^2 is seen and an upper limits on the J/ψ decay branching fractions at the 90% C.L. are given in Table 1.

Table 1: The mass, width, and branching fractions of J/ψ decays into $\{\omega, \phi\}X(1440)$.

$J/\psi \rightarrow \omega X(1440)$ ($X \rightarrow K_S^0 K^+ \pi^- + c.c.$)	$J/\psi \rightarrow \omega X(1440)$ ($X \rightarrow K^+ K^- \pi^0$)
$M = 1437.6 \pm 3.2 \text{ MeV}/c^2$	$M = 1445.9 \pm 5.7 \text{ MeV}/c^2$
$\Gamma = 48.9 \pm 9.0 \text{ MeV}/c^2$	$\Gamma = 34.2 \pm 18.5 \text{ MeV}/c^2$
$B(J/\psi \rightarrow \omega X(1440) \rightarrow \omega K_S^0 K^+ \pi^- + c.c.) = (4.86 \pm 0.69 \pm 0.81) \times 10^{-4}$	
$B(J/\psi \rightarrow \omega X(1440) \rightarrow \omega K^+ K^- \pi^0) = (1.92 \pm 0.57 \pm 0.38) \times 10^{-4}$	
$B(J/\psi \rightarrow \phi X(1440) \rightarrow \phi K_S^0 K^+ \pi^- + c.c.) < 1.93 \times 10^{-5}$ (90% C.L.)	
$B(J/\psi \rightarrow \phi X(1440) \rightarrow \phi K^+ K^- \pi^0) < 1.71 \times 10^{-5}$ (90% C.L.)	

4 New Enhancements

A narrow enhancement is observed in $J/\psi \rightarrow \gamma p\bar{p}$ ¹⁷. Assuming that the $p\bar{p}$ system is in an S-wave resulted in a resonance with mass $M = 1859_{-10-25}^{+3+5}$ MeV/c², width $\Gamma < 30$ MeV/c² (at the 90% C.L.) and product branching fraction $B(J/\psi \rightarrow \gamma X) \cdot B(X \rightarrow p\bar{p}) = (7.0 \pm 0.4(stat)_{-0.8}^{+1.9}(syst)) \times 10^{-5}$. The data not precise enough to determine the angular distribution. According to the theoretical calculation¹⁸, if the X is a bound state of $(p\bar{p})$, the decay channel ($X \rightarrow \eta 4\pi$) is favored over ($X \rightarrow \eta 2\pi, 3\eta$).

The decay channel $J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$ is analyzed using two η' decay modes, $\eta' \rightarrow \pi^+ \pi^- \eta$ and $\eta' \rightarrow \gamma \rho$ ¹⁹. A resonance, the $X(1835)$, is observed with a high statistical significance of 7.7σ in the $\pi^+ \pi^- \eta'$ invariant mass spectrum. From a fit with a Breit-Wigner function, the mass is determined to be $M = 1833.7 \pm 6.1(stat) \pm 2.7(syst)$ MeV/c², the width is $\Gamma = 67.7 \pm 20.3(stat) \pm 7.7(syst)$ MeV/c², and the product branching fraction is $B(J/\psi \rightarrow \gamma X) \cdot B(X \rightarrow \pi^+ \pi^- \eta') = (2.2 \pm 0.4(stat) \pm 0.4(syst)) \times 10^{-4}$. The mass and width of the $X(1835)$ are not compatible with any known meson resonance¹. If we redoing the S -wave BW fit to the $p\bar{p}$ invariant mass spectrum¹⁷ including the zero Isospin, S -wave final-state-interactions (FSI) factor²⁰, yields a mass $M = 1831 \pm 7$ MeV/c² and a width $\Gamma < 153$ MeV/c² (at the 90% C.L.), these values are in good agreement with the mass and width of $X(1835)$ reported here.

In the analysis of $J/\psi \rightarrow \omega p\bar{p}$ ²¹, no significant enhancement near the $p\bar{p}$ mass threshold is observed, and an upper limit of $B(J/\psi \rightarrow \omega X)B(X \rightarrow p\bar{p}) < 1.5 \times 10^{-5}$ is determined at the 95% confidence level.

5 Summary

Using the 58 M J/ψ events sample taken with the BESII detector at the BEPC storage ring, BES experiment provided many interesting results, especially for the study of the lowest glueball candidates, the structure of $\eta(1440)$, and the new enhancement of $X(1835)$, but since the limit of the statistics, the better results (with higher statistics and better accuracy) will be needed for well understanding. The upgraded BEPCII/BESIII will provide a huge J/ψ decay samples for the further analysis.

References

1. W.-M. Yao *et al.*, (Particle Physics Group), *J. Phys. G* **33**, 1 (2006).
2. C. Amsler and F.E. Close, *Phys. Rev. D* **53**, 295 (1996).
3. F.E. Close and A. Kirk, *Eur. Phys. J. C* **21**, 531 (2001).
4. M. Ablikim *et al.* [BES Collab.], *Phys. Lett. B* **642**, 441 (2006).
5. J.Z. Bai *et al.* [BES Collab.], *Phys. Rev. D* **68**, 052003 (2003).
6. M. Ablikim *et al.* [BES Collab.], *Phys. Lett. B* **603**, 138 (2004).
7. M. Ablikim *et al.* [BES Collab.], *Phys. Lett. B* **607**, 243 (2005).
8. H.Y. Cheng, *hep-ph/0609229*.
9. P.H. Baillon *et al.*, *Nuovo Cimento A* **50**, 393 (1967).
10. S. Godfrey and J. Napolitano, *Rev. Mod. Phys.* **71**, 1411 (1999).
11. M.G. Rath *et al.*, *Phys. Rev. D* **40**, 693 (1989).
12. A. Bertin *et al.*, *Phys. Lett. B* **361**, 187 (1995).
13. M. Acciarri *et al.*, *Phys. Lett. B* **501**, 1 (2001).
14. F. Close *et al.*, *Phys. Rev. D* **55**, 5749 (1997).
15. M. Ablikim *et al.* [BES Collab.], *Phys. Lett. B* **594**, 47 (2004).
16. M. Ablikim *et al.* [BES Collab.], *arXiv:0712.1411*.
17. J.Z. Bai *et al.* [BES Collab.], *Phys. Rev. Lett.* **91**, 022001 (2003).

18. G.J. Ding and M.L. Yan, *Phys. Rev. C* **72**, 015208 (2005).
19. M. Ablikim *et al.* [BES Collab.], *Phys. Rev. Lett.* **95**, 262001 (2005).
20. A. Sibirtsev *et al.*, *Phys. Rev. D* **71**, 054010 (2005).
21. M. Ablikim *et al.* [BES Collab.], *Eur. Phys. J. C* **53**, 15 (2008).